

# VON NEUMANN ALGEBRAS OF SOFIC GROUPS WITH $\beta_1^{(2)} = 0$ ARE STRONGLY 1-BOUNDED.

DIMITRI SHLYAKHTENKO

**ABSTRACT.** We show that if  $\Gamma$  is a finitely generated finitely presented sofic group with zero first  $L^2$  Betti number, then the von Neumann algebra  $L(\Gamma)$  is strongly 1-bounded in the sense of Jung. In particular,  $L(\Gamma) \not\cong L(\Lambda)$  if  $\Lambda$  is any group with free entropy dimension  $> 1$ , for example a free group. The key technical result is a short proof of an estimate of Jung [10] using non-microstates entropy techniques.

## 1. INTRODUCTION.

In a series of papers [17, 18], Voiculescu introduced the notion of free entropy dimension, as an analog of Minkowski content in free probability theory. If  $X_1, \dots, X_n \in (M, \tau)$  are a self-adjoint  $n$ -tuple of elements in a tracial von Neumann algebra,  $\delta_0(X_1, \dots, X_n)$  is a measure of “how free they are”. If  $M = W^*(X_1, \dots, X_n)$  satisfies the Connes embedding conjecture and is diffuse,  $1 \leq \delta_0(X_1, \dots, X_n) \leq n$ . The value 1 is achieved e.g. when  $M$  is hyperfinite [8], while a free semicircular  $n$ -tuple generating the free group factor  $L(\mathbb{F}_n)$  has free entropy dimension  $n$  [18].

The number  $\delta_0(X_1, \dots, X_n)$  is an invariant of the (non-closed)  $*$ -algebra generated by  $X_1, \dots, X_n$ . In particular, if  $\Gamma$  is a finitely-generated discrete group, free entropy dimension of any generating set of the group algebra gives us a group invariant,  $\delta_0(\mathbb{C}\Gamma)$  [20]. As it turns out,  $\delta_0(\mathbb{C}\Gamma)$  has a close relationship with the first  $L^2$  Betti number of  $\Gamma$  (we refer the reader to [13] for background on  $L^2$ -Betti numbers). Indeed,  $\delta_0(\mathbb{C}\Gamma) \leq \beta_1^{(2)}(\Gamma) + 1$  for any finitely generated infinite group  $\Gamma$  [5]; and if  $\delta_0$  is replaced by its “non-microstates analog”  $\delta^*$ , then equality holds [14].

Free entropy dimension has found a large variety of applications in von Neumann algebra theory; for example, it was used by Voiculescu to prove that free group factors  $L(\mathbb{F}_n)$  have no Cartan subalgebras [18]. By now, there is a long list of various properties that imply that any generators of a von Neumann algebra  $M$  have free entropy dimension 1; we refer the reader to Voiculescu’s survey [21] and to [9, 7].

In [9] Jung discovered a technical strengthening of the statement that  $\delta_0(X_1, \dots, X_n) = 1$ , called *strong 1-boundedness*, which we will now briefly review.

The free entropy dimension  $\delta_0$  is defined by the formula [18]

$$\delta_0(X_1, \dots, X_n) = n - \limsup_{\epsilon \rightarrow 0} \frac{\chi(X_1 + \epsilon^{\frac{1}{2}} S_1, \dots, X_n + \epsilon^{\frac{1}{2}} S : S_1, \dots, S_n)}{\log \epsilon^{\frac{1}{2}}}$$

where  $\chi$  stands for free entropy and  $S_1, \dots, S_n$  is a free semicircular  $n$ -tuple, free from  $X_1, \dots, X_n$ . The statement that  $\delta_0(X_1, \dots, X_n) \leq 1$  then translates into the estimate  $\chi(X_1 +$

---

Research supported by NSF grant DMS-1500035.

$\epsilon^{\frac{1}{2}}S_1, \dots, X_n + \epsilon^{\frac{1}{2}}S_n : S_1, \dots, S_n) \leq (n-1) \log \epsilon^{\frac{1}{2}} + f(\epsilon)$  with  $\limsup_{\epsilon} |f(\epsilon)/\log \epsilon| = 0$ . Strong 1-boundedness of the set  $X_1, \dots, X_n$  is a strengthening of this inequality: it requires that

$$\chi(X_1 + \epsilon^{\frac{1}{2}}S_1, \dots, X_n + \epsilon^{\frac{1}{2}}S_n : S_1, \dots, S_n) \leq (n-1) \log \epsilon^{\frac{1}{2}} + \text{const}$$

for small  $\epsilon$ .

Jung proved the following amazing result: If  $X = (X_1, \dots, X_n)$  is a strongly 1-bounded set and  $\chi(X_j) > -\infty$  for at least one  $j$ , then any other finite set  $Y = (Y_1, \dots, Y_m) \in W^*(X)$  satisfying  $W^*(Y) = W^*(X)$  is also strongly 1-bounded; thus in particular,  $\delta_0(Y_1, \dots, Y_m) \leq 1$ .

In his follow-up to Jung's work, Hayes [7] improved this statement: if  $X = (X_1, \dots, X_n)$  is strongly 1-bounded and either  $W^*(X)$  is amenable or  $X$  is a non-amenable set, then any  $Y = (Y_1, \dots, Y_m)$  with  $W^*(Y) \cong W^*(X)$  must also be strongly 1-bounded.

We will say that a von Neumann algebra  $M$  is strongly 1-bounded if  $M = W^*(X)$  for some finite strongly 1-bounded set  $X$  which satisfies either of the conditions: (a)  $\chi(X_j) > -\infty$  for some  $j$ , (b)  $W^*(X)$  is amenable, or (c)  $X$  is a non-amenable set. Thus if  $M$  is strongly 1-bounded and  $M = W^*(Y)$  for a finite set  $Y$ , then  $Y$  is strongly 1 bounded and in particular  $\delta_0(Y) \leq 1$ .

Jung's result means that strong 1-boundedness of a von Neumann algebra can be checked on a set of its generators. With few exceptions such as property (T) [12], all known implications stating that some property (presence of a Cartan subalgebra, tensor product decomposition, property  $\Gamma$ , etc.) entails  $\delta_0 = 1$  for any generating set can be upgraded to state that the von Neumann algebra is actually strongly 1-bounded. We refer the reader to [9, 7] and references therein.

The main theorem of this paper states that if a finitely generated finitely presented sofic group  $\Gamma$  satisfies  $\beta_1^{(2)}(\Gamma) = 0$ , then the associated von Neumann algebra is strongly 1-bounded. In particular, any generating set of the von Neumann algebra  $L(\Gamma)$  has free entropy dimension 1. This implies a number of free indecomposability properties of  $L(\Gamma)$ . For example,  $L(\Gamma) \not\cong M * N$  with  $M, N$  diffuse von Neumann algebras; in particular,  $L(\Gamma) \not\cong L(\mathbb{F}_n)$  for any  $n \in [2, +\infty]$ .

To obtain our result, we first give a new proof to a recent result of Jung [10] which allows one to deduce  $\alpha$ -boundedness results for free entropy dimension under the assumption of existence of algebraic relations. Jung's estimate can be considered an improved version of the free entropy dimension estimates from [5, 3] that were previously obtained using non-microstates free entropy methods, but his proof relies on a number of highly technical microstates estimates. Our proof is considerably shorter than Jung's, avoiding much of the difficulty of dealing with matricial microstates. Indeed, we instead produce estimates for the non-microstates Fisher information and non-microstates free entropy and then use results of [2] to deduce the corresponding microstates free entropy inequality and strong boundedness of free entropy dimension. Finally, we apply our estimate to the case of sofic groups.

**Acknowledgments.** I would like to thank Institut Mittag-Leffler and the Hausdorff Institute for Mathematics, where parts of this work were completed, for their support and hospitality. I am also grateful to Ian Charlesworth and Ben Hayes for a number of useful discussions.

## 2. ESTIMATES ON FISHER INFORMATION AND FREE ENTROPY.

**2.1. Preliminaries and notation.** Throughout this section we write  $X = (X_1, \dots, X_n)$  for an  $n$ -tuple of self-adjoint variables in a tracial von Neumann algebra  $(M, \tau)$ . Thus we write  $W^*(X)$  for  $W^*(X_1, \dots, X_n)$ , etc. If  $S = (S_1, \dots, S_n)$  is another  $n$ -tuple we write  $X + \sqrt{\epsilon}S$  for the  $n$ -tuple  $(X_1 + \sqrt{\epsilon}S_1, \dots, X_n + \sqrt{\epsilon}S_n)$ . We also write  $\|X\|_2 = (\sum \|X_j\|_2^2)^{1/2}$ .

We write  $F = (F_1, \dots, F_k)$  for a vector-valued non-commutative polynomial function of  $n$  variables. Here each  $F_j \in \mathbb{C}[t_1, \dots, t_n]$  is a non-commutative polynomial. We write  $\partial F$  for the  $k \times n$  matrix  $(\partial_j F_i)_{ij} \in M_{k \times n}(\mathbb{C}[t_1, \dots, t_n] \otimes \mathbb{C}[t_1, \dots, t_n])$ , where  $\partial_j$  are the non-commutative difference quotient derivations determined by the Leibniz rule and  $\partial_j t_i = \delta_{i=j} 1 \otimes 1$ . We write  $F(X)$ ,  $\partial F(X)$ , etc. when we evaluate these functions by substituting  $X = (X_1, \dots, X_n)$  for  $(t_1, \dots, t_n)$ .

We also view  $\partial F(X)$  as a linear operator in  $M_{k \times n}(W^*(X) \bar{\otimes} W^*(X)^o)$ . It lies in the commutant of the von Neumann algebra  $W^*(X) \bar{\otimes} W^*(X)^o$  acting on  $L^2(W^*(X) \otimes W^*(X)^o)$  by  $(a \otimes b) \cdot (\xi \otimes \eta) = \xi a \otimes b \eta$ . The rank of  $\partial F(X)$  is the Murray-von Neumann dimension (over the algebra  $W^*(X) \bar{\otimes} W^*(X)^o$  of the closure of the image of  $\partial F(X)$ ).

Finally, for a tensor  $Q = a \otimes b$  and an element  $x$  we write  $Q \# x$  for  $axb$ . We use the same notation for the multiplication in the von Neumann algebra  $W^*(X) \bar{\otimes} W^*(X)^o$ . We also use the same notation for  $n$ -tuples and matrices: if  $Q = (Q_{ij})$  is an  $l \times k$  matrix and  $Y = (Y_1, \dots, Y_k)$  is a  $k$ -tuple, we write  $Q \# Y$  for the  $l$ -tuple  $(\sum_{j=1}^k Q_{ij} \# Y_j)_{i=1}^l$ . With this notation we have the following perturbative formula:  $F(X + \sqrt{\epsilon}S) = F(X) + \sqrt{\epsilon} \partial F \# S + O(\epsilon)$ ; it is sufficient to check its validity for monomial  $F$ , which is straightforward.

**2.2. The main estimate on Fisher information  $\Phi^*$ .** In this paper we will denote by  $\log_+(t)$  the function given by  $\log_+(t) = t$  for  $t > 0$  and  $\log_+(0) = 0$ .

**Lemma 2.1.** *Suppose that  $F = (F_1, \dots, F_k) \in \mathbb{C}[t_1, \dots, t_n]^k$  is a vector-valued polynomial non-commutative function of  $n$  variables, and suppose that for some  $X = (X_1, \dots, X_n) \in (M, \tau)$ ,  $F(X) = 0$ . Assume that  $\log_+[\partial F(X)^* \partial F(X)] \in L^1(W^*(X) \otimes W^*(X)^o)$ .*

*Let  $S = (S_1, \dots, S_n)$  be a free semicircular family, free from  $X$ . Then there exist  $\epsilon_0 > 0$  and  $f \in L^1[0, \epsilon_0]$  so that the free Fisher information satisfies the inequality*

$$\Phi^*(X + \sqrt{\epsilon}S) \geq \frac{\text{rank } \partial F(X)}{\epsilon} + f(\epsilon), \quad 0 < \epsilon < \epsilon_0.$$

*Proof.* Denote by  $E_\epsilon$  the conditional expectation from  $W^*(X, S)$  to  $W^*(X + \sqrt{\epsilon}S)$ , and denote by the same letter the extension of  $E_\epsilon$  to  $L^2$ . Then (see [19])

$$\Phi^*(X + \sqrt{\epsilon}S) = \|\xi_\epsilon\|_2^2$$

where

$$\xi_\epsilon = \epsilon^{-1/2} E_\epsilon(S).$$

Since  $F(X) = 0$ , using Taylor expansion we have that  $F(X + \sqrt{\epsilon}S) = \sqrt{\epsilon} \partial F \# S + \zeta_1(\epsilon)$  with  $\|\zeta_1(\epsilon)\|_2 = O(\epsilon)$ . Thus, using contractivity of  $E_\epsilon$  in  $L^2$ , we deduce:

$$\begin{aligned} E_\epsilon(\sqrt{\epsilon} \partial F(X) \# S) &= E_\epsilon(F(X + \sqrt{\epsilon}S)) + E_\epsilon(\zeta_1(\epsilon)) \\ (1) \qquad &= F(X + \sqrt{\epsilon}S) + \zeta_2(\epsilon) \\ &= \sqrt{\epsilon} \partial F(X) \# S + \zeta_3(\epsilon) \end{aligned}$$

with  $\|\zeta_j(\epsilon)\|_2 = O(\epsilon)$ ,  $j = 2, 3$ . Setting  $X_\epsilon = X + \sqrt{\epsilon}S$ , we also have:

$$\begin{aligned} E_\epsilon(\sqrt{\epsilon}\partial F(X)\#S) &= \sqrt{\epsilon}E_\epsilon(\partial F(X_\epsilon)\#S) + \zeta_4(\epsilon) \\ &= \sqrt{\epsilon}\partial F(X_\epsilon)\#E_\epsilon(S) + \zeta_4(\epsilon) \\ &= \sqrt{\epsilon}\partial F(X)\#E_\epsilon(S) + \zeta_5(\epsilon) \end{aligned}$$

where once again  $\|\zeta_j(\epsilon)\|_2 = O(\epsilon)$ ,  $j = 4, 5$ . Combining this with (1) gives

$$\sqrt{\epsilon}\partial F(X)\#E_\epsilon(S) = \sqrt{\epsilon}\partial F(X)\#S + \zeta_6(\epsilon)$$

where  $\|\zeta_6(\epsilon)\|_2 = O(\epsilon)$ . Since  $\xi_\epsilon = \epsilon^{-1/2}E_\epsilon(S)$ , it follows that

$$\partial F(X)\#\xi_\epsilon = \epsilon^{-1/2}\partial F(X)\#S + \zeta'_\epsilon,$$

where  $\|\zeta'_\epsilon\|_2 = O(1)$ .

Let  $E'$  be the orthogonal projection onto the  $L^2$  closure of  $H = \text{span}(W^*(X)SW^*(X)) \subset L^2(W^*(X, S))$ . The map  $Q \mapsto Q\#S$  is an isometric isomorphism of  $L^2(W^*(X) \otimes W^*(X)^o)$  with  $H$  (as is easily verified by direct computation based on the freeness condition).

Since  $H$  is invariant under left and right multiplication by variables from  $X$ ,  $E'$  commutes with  $\partial F(X)\#\cdot$  and thus

$$\partial F(X)\#E'(\xi_\epsilon) = \epsilon^{-1/2}\partial F(X)\#S + \zeta''_\epsilon$$

with  $\|\zeta''_\epsilon\|_2 = O(1)$  and  $\zeta''_\epsilon \in H$ . Applying  $(\partial F)^*\#$  to both sides and denoting  $(\partial F)^*\#(\partial F)$  by  $Q$  finally gives us that there exists an  $0 < \epsilon_0 < 1$  and a constant  $K$  for which

$$Q\#E'(\xi_\epsilon) = \epsilon^{-1/2}Q\#S + \zeta_\epsilon, \quad 0 < \epsilon < \epsilon_0$$

with  $\zeta_\epsilon \in H$ ,  $\|\zeta_\epsilon\|_2 \leq K/2n^{1/2}$ .

Let  $P_\lambda = \chi_{[\lambda, +\infty)}(Q)$  and denote by  $R_\lambda$  the operator  $f_\lambda(Q)$  where  $f(x) = x^{-1}$  for  $x \geq \lambda$  and  $f(x) = 0$  for  $0 \leq x < \lambda$ . Then  $R_\lambda Q = P_\lambda$ .

Since projections are contractive on  $L^2$ , it follows that for any  $\lambda > 0$ ,  $\|P_\lambda\#S\|_2 \leq \|S\|_2 = n^{1/2}$  and furthermore

$$\begin{aligned} \|\xi_\epsilon\|_2^2 &\geq \|P_\lambda E'(\xi_\epsilon)\|_2^2 \\ &= \|R_\lambda\#Q\#E_1(\xi_\epsilon)\|_2^2 \\ &= \|\epsilon^{-1/2}R_\lambda\#Q\#S + R_\lambda\#\zeta_\epsilon\|_2^2 \\ &\geq \epsilon^{-1}\tau(P_\lambda) - 2\epsilon^{-1/2}|\langle P_\lambda\#S, R_\lambda\#\zeta_\epsilon \rangle| \\ &\geq \epsilon^{-1}\tau(P_\lambda) - \epsilon^{-1/2}\lambda^{-1}K. \end{aligned}$$

Let  $r$  be the rank of  $\partial F(X)$  (i.e.,  $r = \lim_{\lambda \rightarrow 0} \tau(P_\lambda)$ , where  $\tau$  denotes the non-normalized trace on  $M_{n \times n}(W^*(X) \bar{\otimes} W^*(X)^o)$ ), and set  $\lambda = \epsilon^{1/4}$ . Let  $\phi(\lambda) = \tau(1 - P_\lambda)$ . Then

$$\begin{aligned} \|\xi_\epsilon\|_2^2 &\geq \epsilon^{-1}(r - \phi(\lambda)) - \epsilon^{-1/2}\lambda^{-1}K \\ &= \frac{r}{\epsilon} - \epsilon^{-1}\phi(\epsilon^{1/4}) - \epsilon^{-3/4}K \\ &= \frac{r}{\epsilon} - \epsilon^{-1}\phi(\epsilon^{1/4}) + f_1(\epsilon) \end{aligned}$$

for some  $f_1 \in L^1[0, \epsilon_0]$ . Thus to conclude the proof, it is enough to show that  $\epsilon^{-1}\phi(\epsilon^{1/4})$  is integrable on  $[0, \epsilon_0]$ .

Let  $d\mu(t)$  be the composition of the trace  $\tau$  with the spectral measure of  $Q$ , so that  $\phi(\lambda) = \int_{0+}^{\lambda} d\mu(t)$ . Let us substitute  $u = \epsilon^{1/4}$  (so  $du = (1/4)\epsilon^{-3/4}d\epsilon$ ,  $u_0 = \epsilon_0^{1/4}$ ) into the integral expression for the  $L^1$  norm of  $\epsilon^{-1}\phi(\epsilon^{1/4})$ :

$$\begin{aligned} \int_{0+}^{\epsilon_0} \frac{1}{\epsilon} \phi(\epsilon^{1/4}) d\epsilon &= \frac{1}{4} \int_{0+}^{u_0} \frac{1}{u} \phi(u) du \\ &= \frac{1}{4} \int_{0+}^{u_0} (\partial_u \log u) \int_0^u d\mu(t) du - \left[ \frac{1}{4} \phi(u) \log u \right]_{0+}^{u_0} \\ &\leq \frac{1}{4} \int_0^{u_0} |\log_+ t| d\mu(t) + \frac{1}{4} \phi(u_0) |\log u_0| + \frac{1}{4} \lim_{\delta \rightarrow 0} |\log \delta| \phi(\delta). \end{aligned}$$

Since by assumption  $\|\log_+ Q\|_1 < \infty$ , the intergral  $\int_0^{u_0} |\log_+ t| d\mu(t)$  is finite. Since if  $\delta < 1$ ,  $|\log \delta| \leq |\log t|$  for  $t \in (0, \delta]$ ,

$$|\log \delta| \phi(\delta) = \int_{0+}^{\delta} |\log \delta| d\mu(t) \leq \int_0^{\delta} |\log_+ t| d\mu(t)$$

we conclude also that  $\lim_{\delta \rightarrow 0} |\log \delta| \phi(\delta)$  is finite. Thus  $\epsilon^{-1}\phi(\epsilon^{1/4}) \in L^1[0, \epsilon_0]$ .  $\square$

*Remark 2.2.* If one drops the assumption that  $\log_+ [\partial F(X)^* \partial F(X)] \in L^1(W^*(X) \otimes W^*(X))$ , the proof of Lemma 2.1 yields the estimate

$$\Phi^*(X + \sqrt{\epsilon}S) \geq \epsilon^{-1} \tau(P_\lambda) - \epsilon^{-1/2} \lambda^{-1} K,$$

where  $\lambda$  is arbitrary. This readily implies that  $\delta^*(X) \leq n - \tau(P_\lambda)$  for all  $\lambda$ , which shows that  $\delta^*(X) \leq n - \text{rank } \partial F(X)$ . This estimate can be easily obtained from [5] by noting that for any finite-rank operator  $T \in FR$ ,

$$0 = [F_j(X), T] = \sum_k [\partial_k F_j(X) \# T, X_k]$$

showing that the dimension of the  $L^2$  closure of the space

$$\{(T_1, \dots, T_n) \in FR^n : \sum_j [T_j, X_j] = 0\}$$

is at least the rank of  $\partial F$  (compare [3]).

*Remark 2.3.* The proof of Lemma 2.1 suggests the following precise expression for the short-time asymptotics of the conjugate variable  $\xi_\epsilon$ :

$$\xi_\epsilon = \epsilon^{-1/2} F \# S + \xi'_\epsilon + O(\epsilon^{1/2})$$

where  $F = (F_{ij})_{i,j=1}^n$  is the projection onto the  $L^2$  closure of the space  $\{(T_1, \dots, T_n) \in FR^n : \sum_j [T_j, X_j] = 0\}$  and  $\xi'_\epsilon = \partial_\epsilon^*(\delta_{ij} 1 \otimes 1 - F)$ . Indeed, the intuition is that  $F$  is the projection onto the space of  $L^2$  cycles, which is perpendicular to the space of  $L^2$  derivations ( $= \ker F$ ). The kernel of  $F$  is a kind of “maximal domain of definition” of  $\partial_{\epsilon=0}^*$ ; then  $\xi'_\epsilon = E_\epsilon(\partial_{\epsilon=0}^*((\delta_{ij} 1 \otimes 1)_{ij} - F))$  is the “bounded part” of  $\xi_\epsilon$  while  $\epsilon^{-1/2} F \# S$  is the “unbounded part” (compare [15]). However, we were unable to prove this exact formula.

### 2.3. Estimates on free entropy.

**Lemma 2.4.** *Under the hypothesis of Lemma 2.1, there exists  $K < \infty$ ,  $\epsilon_0 > 0$ , so that for all  $0 < \epsilon < \epsilon_0$ , the non-microstates free entropy satisfies:*

$$\chi^*(X + \sqrt{\epsilon}S) \leq (\log \epsilon^{\frac{1}{2}})(\text{rank } \partial F(X)) + K$$

*In particular, the microstates entropy also satisfies*

$$\chi(X + \sqrt{\epsilon}S : S) \leq \chi(X + \sqrt{\epsilon}S) \leq (\log \epsilon^{\frac{1}{2}})(\text{rank } \partial F(X)) + K.$$

*Proof.* Let  $\epsilon_0, f$  be as in Lemma 2.1; set  $K' = \|f\|_{L^1[0, \epsilon_0]}$ . By definition [19], up to a universal constant depending only on  $n$ ,

$$\chi^*(X + \sqrt{\epsilon}S) = \frac{1}{2} \int_0^\infty -\Phi^*(X + \sqrt{\epsilon + t}S) + \frac{n}{1+t} dt$$

Thus up to a finite universal constant,

$$\begin{aligned} \chi^*(X + \sqrt{\epsilon}S) &= \frac{1}{2} \int_\epsilon^{\epsilon_0} -\Phi^*(X + \sqrt{t}S) dt \\ &\leq \frac{1}{2} \int_\epsilon^{\epsilon_0} -\frac{\text{rank}(\partial F(X))}{t} + K' dt \\ &= (\log \epsilon^{\frac{1}{2}})(\text{rank } \partial F(X)) - \frac{1}{2} \log \epsilon_0 + K', \end{aligned}$$

where we used the inequality of Lemma 2.1 to pass from the first to the second line.

The inequality for the microstates entropy is due to the general inequalities  $\chi(Z : Y) \leq \chi(Z) \leq \chi^*(Z)$ ; the first is trivial and the second is a deep result of Biane, Capitaine and Guionnet [2].  $\square$

**2.4.  $r$ -boundedness.** The main consequence of our estimates is the following Theorem, originally due to Jung [10, Theorem 6.9]. We give a short alternative proof based on our estimates for non-microstates entropy.

**Theorem 2.5.** *Suppose that  $F = (F_1, \dots, F_k)$  is a vector-valued polynomial non-commutative function of  $n$  variables, and suppose that for some  $X = (X_1, \dots, X_n) \in (M, \tau)$ ,  $F(X) = 0$ . Assume that  $\log_+[\partial F(X)^* \partial F(X)] \in L^1(W^*(X) \otimes W^*(X))$ . Then  $X$  is  $n - \text{rank}(\partial F(X))$  bounded.*

*Proof.* By Lemma 2.4,

$$\limsup_{\epsilon \rightarrow 0} \chi(X + \sqrt{\epsilon}S : S) + \text{rank}(\partial F(X)) |\log \epsilon^{\frac{1}{2}}| \leq K < \infty.$$

Thus by the last bullet point in [9, Corollary 1.4],  $X$  is  $n - \text{rank}(\partial F(X))$ -bounded.  $\square$

*Remark 2.6.* Motivated by [9, Corollary 1.4] one can make the following definition:  $X$  is  $r$ -bounded for  $\delta^*$  (resp.  $\delta^*$ ) if for some  $K$  and all  $0 < \epsilon < \epsilon_0$ ,

$$\chi^*(X + \sqrt{\epsilon}S) \leq (n - r) \log \epsilon^{\frac{1}{2}} + K$$

(respectively,  $\Phi^*(X + \sqrt{\epsilon}S) \geq \epsilon^{-1}(n - r) + \phi(\epsilon)$ ,  $0 < \epsilon < \epsilon_0$  with  $\phi \in L^1[0, \epsilon_0]$ ). Then under the hypothesis of Theorem 2.5,  $X$  is  $n - \text{rank}(\partial F(X))$ -bounded for  $\delta^*$  and  $\delta^*$ .

### 3. APPLICATIONS.

**Lemma 3.1.** *Let  $\Gamma$  be a finitely generated finitely presented group. Then there exists an  $n$ -tuple  $X = (X_1, \dots, X_n)$  of self-adjoint elements in  $\mathbb{Q}\Gamma \subset L\Gamma$  and a vector-valued polynomial function  $F = (F_1, \dots, F_k)$  with  $F_j \in \mathbb{Q}[t_1, \dots, t_n]$  so that  $F(X) = 0$  and moreover*

$$\text{rank } \partial F(X) = n - (\beta_1^{(2)}(\Gamma) + \beta_0^{(2)}(\Gamma) - 1),$$

where  $\beta_j^{(2)}(\Gamma)$  are the  $L^2$ -Betti numbers of  $\Gamma$ .

*Proof.* Let  $g_1, \dots, g_m$  be generators of  $\Gamma$  and let  $n = 2m$ . Consider the following generators  $X_1, \dots, X_n$  for the group algebra  $\mathbb{C}\Gamma$ :  $X_j = g_j + g_j^{-1}$ ,  $X_{m+j} = i(g_j - g_j^{-1})$ ,  $j = 1, \dots, m$ .

Denote by  $h_1, \dots, h_m$  the generators of the free group  $\mathbb{F}_m$ , and let  $Y_j = h_j + h_j^{-1}$ ,  $Y_{n+j} = i(h_j - h_j^{-1})$ ,  $j = 1, \dots, m$  be generators of  $\mathbb{C}\mathbb{F}_m$ . Let  $R_1, \dots, R_p \in \mathbb{F}_m$  be the relations satisfied by the generators  $g_1, \dots, g_m$  of  $\Gamma$ .

Denote by  $t_1, \dots, t_n$  the generators of the algebra  $\mathcal{A} = \mathbb{C}[t_1, \dots, t_n]$  of non-commutative polynomials in  $2n$  variables. Then there is a canonical map from  $\mathcal{A} \rightarrow \mathbb{C}\mathbb{F}_m$  given by  $t_j \mapsto Y_j$ . The algebra  $\mathbb{C}\mathbb{F}_m$  is then the quotient of  $\mathcal{A}$  by the ideal  $J_0$  generated by the relations corresponding to the relations  $h_j^{-1}h_j = h_jh_j^{-1} = 1$  that hold in  $\mathbb{C}\mathbb{F}_m$ ; written in terms of the generators  $t_j$  these relations take the form  $F'_j = (t_j + t_{m+j})(t_j - t_{m+j}) + 4i$ ,  $F''_j = (t_j - t_{m+j})(t_j + t_{m+j}) + 4i$ ,  $j = 1, \dots, m$ .

The relations  $R_j$  can be interpreted as polynomials (with rational coefficients) in  $t_1, \dots, t_n$  by substituting  $\frac{1}{2}(t_j + t_{m+j})$  for  $h_j$  and  $\frac{1}{2i}(t_j - t_{m+j})$  for  $h_j^{-1}$ . Let  $F'''_j = R_j - 1$ ,  $j = 1, \dots, p$ . Let  $J$  be the ideal in  $\mathcal{A}$  generated by  $J_0$  and  $F'''_j$ ,  $j = 1, \dots, p$ . Then  $\mathbb{C}\Gamma$  is precisely the quotient of  $\mathcal{A}$  by  $J$ , the quotient map sending  $t_j$  to  $X_j$ . Let  $F = (F'''_j)_{j=1}^p \sqcup (F''_j)_{j=1}^m \sqcup (F'_j)_{j=1}^m$ , where  $\sqcup$  refers to union of ordered tuples.

Let  $\delta' : \mathcal{A} \rightarrow L^2(\mathbb{C}\Gamma \otimes \mathbb{C}\Gamma^o)$  be a derivation. If for all  $j = 1, \dots, p + 2n$ ,  $\delta'(F_j) = 0$  then for any  $x = aF_jb \in J$ ,

$$\delta'(x) = \delta'(a)(F_j)b + a(F_j)\delta'(b) + a\delta'(F_j)b = 0$$

since  $F_j$  acts by zero on both the right and the left of  $L^2(\mathbb{C}\Gamma \otimes \mathbb{C}\Gamma^o)$  and  $\delta'(F_j) = 0$  by assumption; thus  $\delta'(J) = 0$ . Conversely, if  $\delta'(J) = 0$  then  $\delta(F_j) = 0$  for all  $j$ . It follows that  $\delta'$  descends to a derivation  $\delta : \mathbb{C}\Gamma \rightarrow L^2(\mathbb{C}\Gamma \otimes \mathbb{C}\Gamma^o)$  iff  $\delta'(F_j) = 0$  for all  $j = 1, \dots, p + 2n$ .

Let now  $Q_l \in L^2(\mathbb{C}\Gamma \otimes \mathbb{C}\Gamma^o)$ ,  $l = 1, \dots, n$ . Then there exists a unique derivation  $\delta' : \mathcal{A} \rightarrow L^2(\mathbb{C}\Gamma \otimes \mathbb{C}\Gamma^o)$  so that  $\delta'(t_l) = Q_l$  for all  $l$ . This derivation descends to a derivation  $\delta$  on  $\mathbb{C}\Gamma$  (satisfying  $\delta(X_l) = Q_l$ ) iff for all  $j$ ,

$$0 = \delta'(F_j) = \partial F_j(X) \# Q,$$

i.e.,  $Q \in \ker \partial F(X)$ . It follows that the space of derivations  $Z^1(\mathbb{C}\Gamma, L^2(\mathbb{C}\Gamma \otimes \mathbb{C}\Gamma^o))$  is isomorphic to  $\ker \partial F(X) \subset (L^2(\mathbb{C}\Gamma \otimes \mathbb{C}\Gamma^o))^{\oplus n}$ .

Let  $\alpha$  denote the action of  $L(\Gamma \times \Gamma^o)$  on  $L^2(\mathbb{C}\Gamma \otimes \mathbb{C}\Gamma^o)$  given by  $(g \times h) \cdot (\xi \otimes \eta) = \rho(g)\xi \otimes \rho(h)\eta$ , where  $\rho$  is the right regular representation. If  $\delta : \mathbb{C}\Gamma \rightarrow L^2(\mathbb{C}\Gamma \otimes \mathbb{C}\Gamma^o)$ , then  $\alpha_x \circ \delta$  is again a derivation, for any  $x \in L(\Gamma \times \Gamma^o)$ . It follows that  $L(\Gamma \times \Gamma^o)$  acts on the space of these  $L^2$  derivations; moreover, the isomorphism  $Z^1(\mathbb{C}\Gamma, L^2(\mathbb{C}\Gamma \otimes \mathbb{C}\Gamma^o)) \cong \ker \partial F(X)$  is  $\alpha$ -equivariant.

It is easily seen that

$$\dim_{L(\Gamma \times \Gamma^o)} Z^1(\mathbb{C}\Gamma, L^2(\mathbb{C}\Gamma \otimes \mathbb{C}\Gamma^o)) = \beta_1^{(2)}(\Gamma) - \beta_0^{(2)}(\Gamma) + 1$$

(see e.g. [5, 14]; note that  $\beta_1^{(2)}(\Gamma)$  is the dimension of the space of derivations which are not inner, while  $1 - \beta_0^{(2)}(\Gamma)$  is the dimension of the space of inner derivations).

Putting things together, we obtain that

$$\beta_1^{(2)}(\Gamma) - \beta_0^{(2)}(\Gamma) + 1 = \dim_{L(\Gamma \times \Gamma^o)} \ker \partial F(X) = n - \text{rank } \partial F(X)$$

as claimed. Note that by construction  $F \in \mathbb{Q}[t_1, \dots, t_n]$ .  $\square$

**Theorem 3.2.** *Let  $\Gamma$  be a finitely generated, finitely presented sofic group, and let  $r = \beta_1^{(2)}(\Gamma) - \beta_0^{(2)}(\Gamma) + 1$ . Then there exists a set of generators for  $\mathbb{C}\Gamma$  which is  $r$ -bounded. In particular, if  $|\Gamma| = \infty$  and if  $\beta_1^{(2)}(\Gamma) = 0$ , then  $L(\Gamma)$  is a strongly 1-bounded von Neumann algebra.*

*Proof.* By Lemma 3.1 there exists an  $n$ -tuple  $X = (X_1, \dots, X_n)$  of self-adjoint elements in  $\mathbb{Q}\Gamma \subset L\Gamma$  and a vector-valued polynomial function  $F = (F_1, \dots, F_k)$  with  $F_j \in \mathbb{Q}[t_1, \dots, t_n]$  so that  $F(X) = 0$  and moreover  $\text{rank } \partial F(X) = n - (\beta_1^{(2)}(\Gamma) - \beta_0^{(2)}(\Gamma) + 1)$ .

Furthermore, since  $\partial F(X) \in \mathbb{Q}(\Gamma \times \Gamma^o)$  and  $\Gamma$  (thus also  $\Gamma \times \Gamma^o$ ) is sofic, the determinant conjecture holds [6, 1, 13]. Thus  $\log_+[\partial F(X)^* \partial F(X)] \in L^1(W^*(X) \otimes W^*(X))$ .

We may thus apply Theorem 2.5 to conclude that  $(X_1, \dots, X_n)$  is  $n - \text{rank}(\partial F(X)) = \beta_1^{(2)}(\Gamma) - \beta_0^{(2)}(\Gamma) + 1 = r$  bounded.

Thus if  $\beta_1^{(2)}(\Gamma) = 0$ , it follows that  $r = 1$ .

If  $\Gamma$  is an infinite group,  $\beta_0^{(2)}(\Gamma) = 0$ . If  $\Gamma$  is amenable, it follows [4] that  $L(\Gamma)$  is hyperfinite and thus from the results of [9] we get that  $L(\Gamma)$  is strongly 1-bounded.

Assume that  $\Gamma$  is non-amenable; then any finite generating set of  $\Gamma$  is a non-amenability set (cf. [7]). Since  $\Gamma$  is sofic, its microstates spaces are non-empty. It follows from [7] that  $L(\Gamma)$  is strongly 1-bounded.  $\square$

*Remark 3.3.* The proof of Theorem 3.2 still goes through if we assume that  $\Gamma$  satisfies the determinant conjecture (or, more precisely, that  $\log_+[\partial F(X)^* \partial F(X)] \in L^1(W^*(X) \otimes W^*(X))$  for the specific function  $F$  we are dealing with) and that  $L(\Gamma)$  satisfies the Connes embedding conjecture.

It is an open question whether infinite property (T) von Neumann algebras are always strongly 1-bounded, even in the group case, although it is known that every generating set must have free entropy dimension at most 1 [12]. However, we can settle the case of finitely presented sofic property (T) groups:

**Corollary 3.4.** *Let  $\Gamma$  be an infinite finitely presented sofic property (T) group. Then  $L(\Gamma)$  is strongly 1-bounded.*

*Proof.* Indeed, property (T) implies that  $\beta_1^{(2)}(\Gamma) = 0$  and that the group is finitely generated (see e.g. [13]).  $\square$

As noted in [20, Corollary 4.6], the free entropy dimension  $\delta_0$  is an invariant of the group algebra  $\mathbb{C}\Gamma$  of a discrete group.

**Corollary 3.5.** *Let  $M$  be a finite von Neumann algebra that is not strongly 1-bounded.<sup>1</sup> If  $M \cong L(\Lambda)$  with  $\Lambda$  finitely generated finitely presented sofic, then  $\beta_1^{(2)}(\Lambda) \neq 0$ .*

<sup>1</sup>For example,  $M = W^*(X)$  with  $\delta_0(X) > 1$ , e.g.  $M = L(\Gamma)$  with  $\delta_0(\mathbb{C}\Gamma) > 1$ , e.g.,  $\Gamma = \mathbb{F}_n$ ,  $n \geq 2$ .



*Proof.* If  $\beta_1^{(2)}(\Lambda) = 0$ , Theorem 3.2 would imply that  $L(\Lambda) \cong M$  is strongly 1-bounded. Contradiction.  $\square$

**Conjecture 3.6.** *Let  $\Gamma$  be a finitely presented finitely generated sofic group with  $\beta_1^{(2)}(\Gamma) > 0$ . Then*

- (a)  $L(\Gamma)$  is not strongly one bounded;
- (b)  $\delta_0(\mathbb{C}\Gamma) > 1$ ;
- (c)  $\delta_0(\mathbb{C}\Gamma) = \beta_1^{(2)}(\Gamma) + 1$ .

Note that (c)  $\implies$  (b)  $\implies$  (a) since if  $L(\Gamma)$  were strongly 1-bounded, then  $\delta_0(\mathbb{C}\Gamma)$  would have to be 1. The inequality  $\delta_0(\mathbb{C}\Gamma) \leq \beta_1^{(2)}(\Gamma) + 1$  is known (see [5]) and actually follows from the results of the present paper as well. However, the reverse inequality occurring in statement (c) is only known in a few special cases, e.g.  $\Gamma = \mathbb{F}_n$ , or free products of groups with vanishing first Betti number (possibly with amalgamation over finite or amenable groups) [11, 16]. Nonetheless, it is possible that parts (a) or (b) are more approachable than part (c).

*Remark 3.7.* Together with Corollary 3.5, Conjecture 3.6(a) would imply the following: if  $\Gamma$  and  $\Lambda$  are two finitely generated finitely presented sofic groups and  $L(\Gamma) \cong L(\Lambda)$  then  $\beta_1^{(2)}(\Gamma), \beta_1^{(2)}(\Lambda)$  are either simultaneously zero or simultaneously non-zero. In other words, the vanishing of  $\beta_1^{(2)}$  is a  $W^*$ -equivalence invariant for such groups.

## REFERENCES

- [1] G. Balcı and G. Skandalis. Traces on group  $C^*$ -algebras, sofic groups and Lück's conjecture. Preprint arXiv:1501.05753, 2015.
- [2] P. Biane, M. Capitaine, and A. Guionnet. Large deviation bounds for matrix Brownian motion. *Invent. Math.*, 152(2):433–459, 2003.
- [3] I. Charlesworth and D. Shlyakhtenko. Regularity of polynomials in free variables. Preprint arXiv:1408.0580v2, 2015.
- [4] A. Connes. Classification of injective factors. Cases  $II_1, II_\infty, III_\lambda, \lambda \neq 1$ . *Ann. of Math.*, 104:73–115, 1976.
- [5] A. Connes and D. Shlyakhtenko.  $L^2$ -homology for von Neumann algebras. *J. Reine Angew. Math.*, 586:125–168, 2005.
- [6] G. Elek and E. Szabó. Hyperlinearity, essentially free actions and  $\ell_2$ -invariants. the sofic property. *Mathematische Annalen*, 332(2):421–441, 2005.
- [7] B. Hayes. 1-bounded entropy and regularity problems in von Neumann algebras. Preprint arXiv:1505.06682, 2015.
- [8] K. Jung. A hyperfinite inequality for free entropy dimension. *Proc. Amer. Math. Soc.*, 134(7):2099–2108 (electronic), 2006.
- [9] K. Jung. Strongly 1-bounded von Neumann algebras. *Geom. Funct. Anal.*, 17(4):1180–1200, 2007.
- [10] K. Jung. The rank theorem and  $L^2$ -invariants in free entropy: global upper bounds. Preprint arXiv:1602.04726, 2016.
- [11] N. Brown, K. Dykema and K. Jung. Free entropy dimension in amalgamated free products. *Proc. London Math. Soc.* 97(3): 339D367, 2008.
- [12] K. Jung and D. Shlyakhtenko. Any generating set of an arbitrary property  $T$  von Neumann algebra has free entropy dimension  $\leq 1$ . *J. Noncommut. Geom.*, 1(2):271–279, 2007.
- [13] W. Lück.  $L^2$ -invariants: theory and applications to geometry and  $K$ -theory, volume 44 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2002.

- [14] I. Mineyev and D. Shlyakhtenko. Non-microstates free entropy dimension for groups. *Geom. Func. Anal.*, 15:476–490, 2005.
- [15] D. Shlyakhtenko. Some estimates for non-microstates free entropy dimension, with applications to  $q$ -semicircular families. *IMRN*, 51:2757–2772, 2004.
- [16] D. Shlyakhtenko. Lower estimates on microstates free entropy dimension. *Analysis and PDE*, 2:119–146, 2009.
- [17] D.-V. Voiculescu. The analogues of entropy and of Fisher’s information measure in free probability theory II. *Invent. Math.*, 118:411–440, 1994.
- [18] D.-V. Voiculescu. The analogues of entropy and of Fisher’s information measure in free probability theory, III. *Geometric and Functional Analysis*, 6:172–199, 1996.
- [19] D.-V. Voiculescu. The analogues of entropy and of Fisher’s information measure in free probability, V. *Invent. Math.*, 132:189–227, 1998.
- [20] D.-V. Voiculescu. A strengthened asymptotic freeness result for random matrices with applications to free entropy. *IMRN*, 1:41 – 64, 1998.
- [21] D.-V. Voiculescu. Free entropy. *Bull. London Math. Soc.*, 34(3):257–278, 2002.

DEPARTMENT OF MATHEMATICS, UCLA, LOS ANGELES, CA 90095, USA  
*E-mail address:* shlyakht@math.ucla.edu